Cervical Cancer Can Be Ended in the US and Around the World

Each year in the United States, about 11,500 new cases of cervical cancer are diagnosed. Around 4,000 women die of this cancer.

But cervical cancer can be easily prevented. Once diagnosed, it is **one of the most treatable forms of cancer.**

January Is Cervical Cancer Awareness Month

How does one protect against cervical cancer?

Cervical cancer develops in a woman's cervix, the lower part of the uterus. Early-stage cervical cancer may have no symptoms. Almost all cases (99%) of the cancer are caused by the human papillomavirus (HPV), a very common virus that spreads through skin-to-skin contact. The HPV vaccine protects girls and women from cervical cancer.

Getting vaccinated and being screened regularly protects you against cervical cancer.

Who should get vaccinated?

All females aged 9–14 years should get the HPV vaccination. The CDC recommends getting the HPV vaccine at age 11 or 12 as it produces a stronger immune response during the preteen years.

Vaccination is also recommended through age 26.

After age 30, women should have **regular screenings** for other types of cervical cancer that are not prevented by the vaccine.

How does one pay for the vaccine?

- Most insurance companies cover the cost of vaccination.
- The Vaccines for Children (VFC) Program is a federal program that offers vaccines at no cost for eligible children aged 18 and under.
- Other options include checking with a local Planned Parenthood clinic or health department.

Get informed

Educate yourself and other women about cervical cancer and HPV, causing 99% of the cases.

Get screened

regular screenings can detect any cervical changes and can protect you from developing cancer.

Get vaccinated

Girls **between 9 to 14 years** should receive one dose of **HPV vaccine.**

World Health Organization

healthcare education project

Healthcare • Education • Action!

498 7TH AVENUE, 5TH FLOOR, NEW YORK, NY 10018 • TELEPHONE: (646) 473-8481 • WEBSITE: WWW.HEALTHCAREEDUCATIONPROJECT.ORG

SCAN HERE FOR MORE RESOURCES

El Cáncer Cervical Puede Ser Erradicado en los Estados Unidos y Alrededor del Mundo

Cada año en los Estados Unidos, se diagnostican aproximadamente 11,500 nuevos casos de cáncer cervical. Alrededor de 4,000 mujeres mueren de este cáncer.

Pero el cáncer cervical se puede prevenir fácilmente. Una vez diagnosticado, **este cáncer es uno de los más tratables.**

Enero es el Mes de Concientización del Cáncer Cervical, También Conocido Como Cáncer de Cuello Uterino

Còmo se protege contra el cáncer cervical?

El cáncer cervical se desarrolla en el cuello uterino de una mujer, en la parte inferior. El cáncer cervical en etapa temprana puede no tener síntomas. Casi todos los casos (99%) del cáncer son causados por el virus del papiloma humano (HPV), un virus muy común que se propaga a través del contacto piel con piel. La vacuna contra el HPV protege a las niñas y mujeres del cáncer cervical.

Vacunarse y hacerse pruebas de detección regularmente le protege contra el cáncer cervical.

Quièn debería vacunarse?

Todas las mujeres de 9 a 14 años deberían vacunarse contra el HPV. Los CDC (Centros para el Control y la Prevención de Enfermedades) recomiendan recibir la vacuna contra el HPV (virus del Papiloma Humano).

También se recomienda la vacunación hasta los 26 años.

Después de los 30 años, las mujeres deben someterse a **exámenes regulares** de detección de otros tipos de cáncer cervical que no se previenen con la vacuna.

¿Cómo se paga la vacuna?

- La mayoría de las compañías de seguros cubren el costo de la vacunación.
- El Programa de Vacunas Para Niños (VFC) es un programa federal que ofrece vacunas sin costo para niños elegibles de 18 años o menos.
- Otras opciones incluyen consultar con una clínica local de Planned Parenthood (Paternidad Responsable) o con el departamento de salud.

Infórmate

Edúquese a sí misma y a otras mujeres sobre el cáncer cervical y el HPV, causa del 99% de los casos.

Ser examinado

Los exámenes regulares pueden detectar cualquier cambio cervical y pueden protegerte del inicio de cáncer cervical llamado también cáncer de cuello uterino.

Vacunarse

Las niñas entre 9 y 14 años deben recibir una dosis de la vacuna contra el HPV.

World Health Organization

healthcare education project

Healthcare • Education • Action!

498 7TH AVENUE, 5TH FLOOR, NEW YORK, NY 10018 • TELÉFONO: (646) 473-8481 • SITIO WEB: WWW.HEALTHCAREEDUCATIONPROJECT.ORG

ESCANEE AQUÍ PARA OBTENER MÁS RECURSOS